

RESOLVED ELECTRON*

Wojciech SŁOMIŃSKI and Jerzy SZWED

Institute of Computer Science, Jagellonian University, Reymonta 4, 30-059 Kraków, Poland

Advantages of introducing the electron structure function in electron induced processes are demonstrated. At present energies the same experiment gives more precise description of the electron than photon structure. The momentum scales entering the process are better controlled. At very high momenta probabilistic (partonic) interpretation can be preserved despite strong γ - Z interference. The “virtual photon” structure can also be reformulated in terms of more physical (real) electron variables.

Theoretical framework which allows to calculate the photon structure is known since long [1]. It appears as perturbative QCD contribution (resolved photon), in addition to the modeled Vector Meson Dominance (VDM) and pointlike (direct) terms. To measure this photonic structure, experiments [2] use the electron (or positron) beam as a source of photons. Despite precise measurement the data are difficult to extract. The problem is displayed in Fig. 1a. The tagged (upper) electron emits a probing photon whereas the untagged (lower) goes nearly along the beam, emitting the target photon. The large scale, Q^2 is determined by the tagged electron:

$$Q^2 = -(k - k')^2 = 2EE_{\text{tag}}(1 - \cos \theta_{\text{tag}}), \quad (1)$$

where E is the initial electron energy and E_{tag} and θ_{tag} are the energy and polar angle of the measured electron. The anti-tagging condition (if present) requires the virtuality of the target photon to be less than P^2 :

$$-(p - p')^2 \equiv P_\gamma^2 \leq P^2.$$

This photon is clearly not a beam particle and has the energy diffused according to the equivalent photon (Weizsäcker-Williams [3]) spectrum f_γ^e . The measured cross-section for the production of a hadronic system X , expressed in terms of the photon structure functions F_2^γ and F_L^γ reads:

$$\begin{aligned} \frac{d^2 \sigma_{ee \rightarrow eX}}{dz dQ^2} &= \frac{2\pi\alpha_{\text{em}}^2}{x^2 Q^4} \\ &\times [(1 + (1 - y)^2) F_2^\gamma(x, Q^2, P^2) - y^2 F_L^\gamma(x, Q^2, P^2)] \\ &\times f_\gamma^e(z/x, P^2) dx \end{aligned} \quad (2)$$

where

$$y = 1 - (E_{\text{tag}}/E) \cos^2(\theta_{\text{tag}}/2)$$

and $x(z)$ are fractions of parton momentum with respect to the photon (electron).

Three remarks are important for further considerations. First, the splitting of the process into a distribution of photons inside electron f_γ^e and that of partons

inside the photon F_2^γ is an approximation. The optimal form of the equivalent photon formula is still being discussed [4]. Even if most of experimental groups choose the same formula, one should keep in mind that the definition of the photon structure function depends on this convention.

Second, in order to fix x , one is forced to measure — in addition to the tagged electron — the hadronic momenta. In fact,

$$x = \frac{Q^2}{Q^2 + P_\gamma^2 + W^2} \approx \frac{Q^2}{Q^2 + W^2},$$

where W is the invariant mass of the produced hadronic system X . Its determination is more difficult than other (tagged electron) variables.

Third, the target photon is always off-shell. Although the equivalent photon distribution is peaked at minimum (nearly zero) virtuality, treating the photon as real is another approximation. One should keep in mind that the measured photon structure function depends on x , Q^2 and P^2 (we keep the minimum virtuality fixed but in general it is still another variable).




FIG. 1. Deep inelastic scattering on a photon (a) and electron (b) target

The necessity of unfolding the photon structure function from the cross-section (2) and the uncertainty in the determination of the x variable are sources of biggest errors in the analysis. The data are indirectly biased by theoretical assumptions and, in addition, the selection cuts put on the hadronic mass W^2 reduce the number of accepted events.

Most of the above problems can be avoided when we introduce the structure function of the electron (Fig. 1b). To see how it works let us first write the cross-section (2), this time in terms of the electron structure functions F_2^e and F_L^e :

$$\frac{d^2\sigma_{ee\rightarrow eX}}{dzdQ^2} = \frac{2\pi\alpha_{\text{em}}^2}{zQ^4} [(1 + (1-y)^2)F_2^e(z, Q^2, P^2) - y^2 F_L^e(z, Q^2, P^2)]. \quad (3)$$

The structure function $F_2^e(z, Q^2, P^2)$ which dominates the cross-section at small y , has simple partonic interpretation:

$$F_2^e(z, Q^2, P^2) = z \sum_i e_{q_i}^2 q_i(z, Q^2, P^2)$$

where e_{q_i} and q_i are the i -th quark fractional charge and density. No unfolding procedure is necessary to obtain F_2^e , and its argument z — the parton momentum fraction with respect to the electron — is measured, as in the standard deep inelastic scattering, with the tagged electron variables only:

$$z = \frac{Q^2}{2pq} = \frac{\sin^2(\theta_{\text{tag}}/2)}{E/E_{\text{tag}} - \cos^2(\theta_{\text{tag}}/2)}.$$

There is no need to reconstruct the hadronic mass W . All these features cause that the same experiment can produce more precise data. What is most important — the electron structure function contains the same information about QCD as the photon one and, as we point out below, it is known theoretically with at least the same accuracy. Moreover, it allows to avoid problems which arise in the photon structure function at very high energies.

The construction of the QCD electron structure function can be presented in two steps. First we calculate the splitting function of the electron into a quark/anti-quark $P_{q/\bar{q}}^e(z, P^2)$ (Fig. 2). As a high Q^2 probe we use in this calculation the gluon rather than the photon, because it couples to the QCD partons only. Note that we allow for the exchange of Z and W bosons in addition to the photon. The result for a quark reads (for anti-quark: exchange Φ_+ with Φ_- and replace δ_{qd} with $\delta_{\bar{q}\bar{u}}$):

$$P_q^e(z, P^2) = \frac{3\alpha_{\text{em}}}{4\pi} \left\{ 2e_q^2 [\Phi_+(z) + \Phi_-(z)] \log \mu_0 + \tan^4 \theta_W [(e_q^2 + z_q^2 \rho^2) \Phi_+(z) + (e_q^2 \rho^2 + z_q^2) \Phi_-(z)] \log \mu_Z + 2e_q \tan^2 \theta_W [(e_q + z_q \rho) \Phi_+(z) - (e_q \rho + z_q) \Phi_-(z)] \log \mu_Z + (1 + \rho)^2 \Phi_+(z) \delta_{qd} \log \mu_W \right\} \quad (4)$$

where

$$\Phi_+(z) = \frac{1-z}{3z} (2 + 11z + 2z^2) + 2(1+z) \log z, \quad (5)$$

$$\Phi_-(z) = \frac{2(1-z)^3}{3z} \quad (6)$$

$\rho = 1/2 \sin^2 \theta_W - 1$, $\mu_0 = P^2/m_e^2$, $\mu_B = P^2/M_B^2 + 1$, $z_q = T_3^q/\sin^2 \theta_W - e_q$, with e_q , T_3^q , M_B , m_e and θ_W being the quark charge, its 3rd weak isospin component, weak boson (Z or W) mass, electron mass and the Weinberg angle. One sees that at present Q^2 the photon contribution (first term proportional to $\log \mu_0$) dominates. At very high momenta one should expect the other terms to contribute, in particular the γ - Z interference enters with the same logarithm as the Z -term itself.




FIG. 2. The Feynman diagrams contributing to the electron $\rightarrow$ quark/anti-quark splitting functions

In the second step we construct the Q^2 -evolution equations for the quark and gluon densities inside the electron $q(z, Q^2, P^2)$ and $G(z, Q^2, P^2)$. Introducing $t = \log(Q^2/\Lambda_{\text{QCD}}^2)$ and $t_1 = \log(P^2/\Lambda_{\text{QCD}}^2)$ and remembering that there is no direct coupling of the electroweak sector to gluons we can write [5]:

$$\frac{dq_j(t, t_1)}{dt} = \frac{\alpha_{\text{em}}}{2\pi} P_{q_j}^e(t_1) + \frac{\alpha(t)}{2\pi} \sum_k P_{q_j}^{qk} \otimes q_k(t, t_1) + P_{q_j}^G \otimes G(t, t_1), \quad (7)$$

$$\frac{dG(t, t_1)}{dt} = \frac{\alpha(t)}{2\pi} \sum_k P_G^{qk} \otimes q_k(t, t_1) + P_G^G \otimes G(t, t_1). \quad (8)$$

The sign $\otimes$ stands for the convolution $f \otimes g = \int_z^1 \frac{dx}{x} f(z/x)g(x)$ and explicit z dependence has been suppressed. $\alpha(t)$ is the QCD running coupling constant.

The momentum scales require some attention. The QCD evolution, started by the $q\bar{q}$ pair, is governed by Λ_{QCD} . The large ‘probing scale’ defined for e^+e^- scattering in Eq.(1), is chosen in other reactions to be large transverse momentum (e.g. of jets) or produced heavy mass (e.g. heavy quark or Higgs boson). The maximum virtuality P^2 defines the kinematical range which is called photo- (or more general boson-) production. It is often set by experiment. E.g. in e^+e^- or ep scattering it can be chosen by the untagging condition. But one may also take into account larger virtualities up to $Q^2 \lesssim P^2$. In the case of the W boson contribution no such limit can

be imposed experimentally at all and P^2 varies up to its kinematical limit:

$$P_{\max}^2 = \frac{x+1-z}{x} Q^2.$$

In the asymptotic t region ($t \rightarrow \infty$) we can take [6]

$$t_1 = (1-a)\hat{t}_1 + at$$

where $\hat{t}_1$ is constant and $0 \leq a \leq 1$. Let us look closer at the two limiting cases $a = 0$ and $a = 1$. In the first one the inhomogeneous term in the evolution equations (7) does not depend on t . The asymptotic solutions have the form:

$$\begin{aligned} q(z, t) &\simeq \left(\frac{\alpha_{\text{em}}}{2\pi}\right)^2 q^{\text{as}}(z) t_1 t, \\ G(z, t) &\simeq \left(\frac{\alpha_{\text{em}}}{2\pi}\right)^2 G^{\text{as}}(z) t_1 t \end{aligned} \quad (9)$$

with $q^{\text{as}}(z)$ and $G^{\text{as}}(z)$ being given by known, t -independent integral equations [5,6].

Their numerical solutions are shown in Fig. 3. As expected in the asymptotic region all bosons contribute. What is interesting, the γ - Z interference term enters with strength comparable to the contribution of the Z boson itself. This means that the notion of separate equivalent bosons breaks down at very high momenta. The electron structure function takes correctly into account interference effects, preserving at the same time probabilistic (partonic) interpretation.

One should keep in mind that only in the asymptotic region all logarithms entering the splitting function P_q^e are equal. At lower energies the photon contribution (proportional to $\log \mu_0$) dominates.





FIG. 3. Asymptotic quark and gluon distributions — solid line. The other lines show different contributions as labeled

The ‘inclusive’ (no tagging) case when $a = 1$ was considered in Ref. [5,6]. After quite non-trivial procedure the evolution equations look formally identical to Eqs.(7) with t_1 replaced by t . In particular the splitting function P_q^e is now proportional to t . The solutions again have the form (9) with t_1 replaced by t (the parton densities grow thus as t^2), but the resulting z -dependence is different from the previous case. The standard convolution of equivalent photons with the photon structure (as in case $a = 0$) is no more valid.

The comparison of the two cases is shown for the electron structure function $F_2^e(z, Q^2, P^2)$ in Fig. 4. Two upper curves are the asymptotic result with all electroweak bosons taken into account. We also show the photon contribution alone to get a feeling what is the effect at presently accessible momenta. One sees an extra suppression in the ‘inclusive’ case (when $t_1 \sim t$).

Let us add a few final remarks. First concerns the study of the virtual photon structure [7]. The analysis can be reformulated in terms of the P^2 dependence of the electron structure function. Studying a real, convention independent object is first advantage. Another one is the fact that at very high virtualities the Z admixture and the γ - Z interference are properly taken into account.




FIG. 4. $F_2^e(z, Q^2, P^2)/(\frac{\alpha_{em}}{2\pi})^2 t_1 t$ for $P^2 \ll Q^2$ — solid line, and for $P^2 \approx Q^2$ — broken line. The contributions from γ, Z, W and from γ alone are labeled

Second, the calculation discussed above has been done in leading logarithmic approximation. Corrections to this picture are well defined. In particular one does not require factorization of the electron induced process into the $e\text{-}\gamma$ (Weizsäcker-Williams) and γ -QCD parts. The process initializing the $q\bar{q}$ cascade can be calculated exactly (in given order of α_{em}).

Third, the extraction of the electron structure function consists in the reanalysis of the collected e^+e^- data which served for the photon. A phenomenological parametrisation describing the data, with the predicted Q^2 and P^2 dependence, could be then constructed (including the VDM and direct contributions, see also [8]). This parametrisation would serve then as the basis for a Monte-Carlo generator simulating quark production in lepton induced processes. This single generator replaces in fact two others, presently used [9].

As a side-remark we comment on the QED structure function of the photon. It is extracted from the process $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ by dividing out the (approximate) equivalent photon distribution. The use of the QED electron structure function avoids this problem. The exactly known (in given order of α_{em}) electron structure function can be compared directly with the electron data.

Finally, the photon structure function has been recently measured [10] in two jet production at HERA. Again the extraction of the x variable is difficult. In addition to jets, one has to measure essentially the whole hadronic system in order to obtain the photon energy. The data, when presented in terms of the electron structure function, require only measurement of the two jets. As in the e^+e^- case the resulting data should be more precise.

To summarize, we propose to look at the electron as surrounded by a QCD cloud of quarks and gluons (in order α_{em}^2), very much like it is surrounded by a QED cloud of equivalent photons (in order α_{em}). We argue that the use of the electron structure function in electron induced

processes has important advantages over the photon one. Experimentally it leads to more precise, convention independent data. Theoretically it allows for more careful treatment of all variables. It also takes into account all electroweak gauge boson contributions, including their interference, which will be important in the next generation of e^+e^- colliders.

The authors would like to thank Maria Krawczyk, Aharon Levy and Jacek Turnau for discussions. The hospitality of the DESY Theory Group is also acknowledged.

* Work supported by the Polish State Committee for Scientific Research (grant No. 2 P03B 081 09) and the Volkswagen Foundation.

- [1] E. Witten, Nucl. Phys. **B120**, 189 (1977);
C.H. Llewellyn-Smith, Phys. Lett. **79B**, 83 (1978);
R.J. DeWitt et al., Phys. Rev. **D19**, 2046 (1979);
T.F. Walsh and P. Zerwas, Phys. Lett. **36B**, 195 (1973);
R.L. Kingsley, Nucl. Phys. **B 60**, 45 (1973).
- [2] H.J. Behrend et al., CELLO Collaboration, Phys. Lett. **126B**, 391 (1983); Ch. Berger et al., PLUTO Collaboration, Phys. Lett. **142B**, 111 (1984); Nucl. Phys. **B281**, 365 (1987); W. Bartel et al., JADE Collaboration, Zeit. f. Phys. **C24**, 231 (1984); M. Althoff et al., TASSO Collaboration, Zeit. f. Phys. **C31**, 527 (1986); H. Aihara et al., TPC/2 γ Collaboration, Zeit. f. Phys. **C34**, 1 (1987); Phys. Rev. Lett. **58**, 97 (1987); T. Sasaki et al., AMY Collaboration, Phys. Lett. **252B**, 491 (1990); R. Akers et al., OPAL Collaboration, Zeit. f. Phys. **C61**, 199 (1994); S.K. Sahu et al., AMY Collaboration, Phys. Lett. **B346**, 208 (1995); P. Abreu et al., DELPHI Collaboration, Zeit. f. Phys. **C69**, 223 (1996); K. Ackerstaff et al., OPAL Collaboration, Zeit. f. Phys. **C74**, 33 (1997); Phys. Lett. **B411**, 387 (1997); *ibid.* **B412**, 225 (1997).
- [3] C.F. von Weizsäcker Z. Phys. **88**, 612 (1934); E.J. Williams, Phys. Rev. **45**, 729 (1934).
- [4] A.C. Bawa and W.J. Stirling, Journal of Physics, **G15**, 1339 (1989); S. Frixione, M.L. Magnano, P. Nason and G. Ridolfi, Phys. Lett. **B319**, 339 (1993).
- [5] W. Słomiński and J. Szwed, Phys. Lett. **B387**, 861 (1996); Acta Phys. Polon. **B27** 1887 (1996).
- [6] W. Słomiński, Jagellonian University preprint 1998.
- [7] T. Uematsu and T.F. Walsh, Phys. Lett. **B101**, 263 (1981); F.M. Borzumati and G.A. Schuler, Zeit. f. Phys. **C58**, 139 (1993).
- [8] M. Drees and R.M. Godbole, Phys. Rev. **D50**, 3124 (1994).
- [9] see e.g. CERN 96-01 Physics at LEP2, ed. G. Altarelli, T. Sjöstrand and F. Zwirner, (1996) Vol. 2, p. 188.
- [10] H1 collaboration, C. Adloff et al., Eur. Phys. J. **C1**, 97 (1998), ZEUS collaboration, J. Breitweg et al., Eur. Phys. J. **C1**, 109 (1998).